

BI AK85-

For

BES
LIBRARY

Rec- 12123185
AW

FINAL TECHNICAL REPORTS FOR RESEARCH UNIT 643

EVALUATION OF THE EFFECTS OF OIL DEVELOPMENT
ON THE COMMERCIAL FISHERIES
IN THE EASTERN BERING SEA

T. Laevastu and R. Marasco
Project Leaders, and

N. Bax^{1/}, R. Fredin^{2/}, F. Fukuhara^{2/}, A. Gallagher, T. Honkalehto^{1/},
J. Ingraham, P. Livingston, R. Miyahara, and N. Pola

Resource Ecology and Fisheries Management Division
Northwest and Alaska Fisheries Center, Seattle, WA 98115
and

1/ Compass Systems Inc.
2/ Natural Resource Consultants

November 1985

Minerals Management Service/National Oceanic and Atmospheric Administration
Interagency Agreement, IA - 14-12-001-30019

ESTIMATED IMPACTS OF HYPOTHETICAL OIL SPILL ACCIDENTS OFF
PORT **MOLLER**, PORT **HEIDEN** AND CAPE **NEWENHAM**
ON EASTERN BERING SEA YELLOWFIN SOLE

By

Francis M. **Fukuhara**

Institute for Marine Studies
University of Washington
Seattle, Washington 98125

and

Natural Resources Consultants
4055 21st Avenue West
Seattle, Washington 98199

Report to RU 643
National Ocean Service
Office of Oceanography and Marine Service
Anchorage, **Alaska**

Resource Ecology and Ecosystem Simulation Task
Resource Ecology and Fisheries Management Division
Northwest and Alaska Fisheries Center
Building 4, BIN **C15700**
7600 Sand Point Way **N.E.**
Seattle, WA 98115



Northwest and
Alaska Fisheries
Center

National Marine
Fisheries Service

U. S. DEPARTMENT OF COMMERCE

NWAFRC PROCESSED REPORT 85-15

Estimated Impacts
of Hypothetical Oil Spill Accidents
off
Port Moller, Port Heiden
and Cape Newenham
on Eastern Bering Sea
Yellowfin Sole

July 1985

ACKNOWLEDGEMENTS

"This study was funded wholly by the Minerals Management Service, Department of the Interior, through an **Interagency** Agreement with the National Oceanic and Atmospheric Administration, Department of Commerce, as part of the **Alaska** Outer Continental Shelf **Environmental** Assessment Program."

This is an advance copy of the "**Final Report.**" Minor editorial changes have been made to the report with concurrence of the National Oceanic and Atmospheric Administration Contracting Officer's **Technical Representative (COTR)**. The "**Final Report**" is being processed through the NOAA-Outer Continental Shelf **Environmental** Assessment Program editing and publishing system and should be available in the near future.

LIST OF FIGURES

	<u>Page</u>
Figure 1. Overall distribution and areas of commercial fishing for yellowfin sole.	3
Figure 2. Seasonal changes in relative abundance of yellowfin sole as shown by NWAF C trawl surveys.	5
Figure 3. Distribution and relative abundance by weight of yellowfin sole in the eastern Bering Sea.	6
Figure 4. Seasonal changes in Japanese fishing grounds for yellowfin sole, December 1967 to October 1968 (Wakabayashi 1974).	8
Figure 5. National Marine Fisheries Service trawl survey subareas showing locations of hypothetical oil spill sites off Port Moller , Port Heiden and Cape Newenham .	10
Figure 6. Percentage distribution of biomass of yellowfin sole in subareas of the NWAF C eastern Bering Sea trawl survey.	14
Figure 7. Distribution of yellowfin sole eggs as shown by plankton surveys.	15
Figure 8. Distribution of yellowfin sole by size group, August- October 1975.	17
Figure 9. Von Bertalanffy growth curves for yellowfin sole taken during spring 1976 (Bakkala and Smith 1978).	23
Figure 10. Age composition of yellowfin sole of the eastern Bering Sea as shown by data from trawl surveys of the Northwest and Alaska Fisheries Center and by U.S. observer data from the commercial fishery.	31
Figure 11. Relative abundance (CPUE) for Japanese pair trawlers and NWAF C trawl surveys.	35
Figure 12. Time-area restrictions applicable to non-U.S. groundfish fisheries in the eastern Bering Sea and Aleutian Islands regions.	43

LIST OF TABLES

	<u>Page</u>
Table 1. Biomass estimates of eastern Bering Sea yellowfin sole from research vessel surveys 1975-1984.	11
Table 2. Estimated biomass of yellowfin sole in subareas of the eastern Bering Sea 1975-83.	13
Table 3. Ratio of females to males by regions and seasons.	19
Table 4. Ratio of females to males in all areas, Subarea 1 and Subarea 4S.	18
Table 5. Fecundity of yellowfin sole in the eastern Bering Sea.	20
Table 6. Estimated number of eggs produced by yellowfin sole in Subareas 1 and 4S in 1983.	21
Table 7. Von Bertalanffy growth parameters for yellowfin sole of eastern Bering Sea.	24
Table 8. Six year means of observed lengths and calculated weights at age for yellowfin sole of the eastern Bering Sea from NWAF survey data of 1973-78.	26
Table 9. Stomach contents (in grams) by size group of yellowfin sole collected in eastern Bering Sea in 1970 (Wakabayashi 1974).	28
Table 10. Species showing close association with yellowfin sole as indicated by recurrent group analysis.	30
Table 11. Estimated number of yellowfin sole (billions of fish) in the eastern Bering Sea, 1959-81, based on cohort analysis.	33
Table 12. Estimated biomass (in 1000 t) of yellowfin sole in the eastern Bering Sea, 1959-81, based on cohort analysis.	34
Table 13. Annual catches of yellowfin sole in the eastern Bering Sea (east of long. 180° and north of lat. 54°) in metric tons.	37
Table 14. Yellowfin sole catch by month, 1979-84.	39
Table 15. Total allowable catch (TAC) and its allocation under the FCMA .	41
Table 16. Forecast of yellowfin sole abundance in the eastern Bering Sea, 1982-89, with constant catches of 214,500 t, natural mortality = 0.12, and recruitment at the lower and higher estimates for 1959-81.	45

Table 17.	Petroleum hydrocarbon uptake expressed as counts per minute (cpm) of radioactive tracer, embryo mortality, hatching success, total length of newly-hatched (mm), and malformed newly-hatched California halibut larvae following 72 hours of embryonic test solution exposure. All starred (*) elements were statistically significantly different from control values.	47
Table 18.	Summary of effects of petroleum hydrocarbons on flatfish.	54
Table 19.	Hypothetical oil spill scenarios.	56
Table 20.	Estimated concentrations and area contaminated by simulated oil spills at Port Heiden , Port Moller and off Cape Newenham.	59
Table 21.	Estimated percentage of mortality from acute toxicity in yellowfin sole in the accident scenarios at Port Moller , Port Heiden and Cape Newenham by life history group and quarter.	61